arXiv:math/0311137v2 [math.AG] 23 Jan 2004

NON-SUPERSINGULAR HYPERELLIPTIC JACOBIANS

YURI G. ZARHIN

Abstract. In his previous papers [25, 26, 28] the author proved that in characteristic $\neq 2$ the jacobian $J(C)$ of a hyperelliptic curve $C : y^2 = f(x)$ has only trivial endomorphisms over an algebraic closure K_a of the ground field K if the Galois group $\mathrm{Gal}(f)$ of the irreducible polynomial $f(x) \in K[x]$ is either the symmetric group $\mathbf{S}_n$ or the alternating group $\mathbf{A}_n$. Here $n \geq 9$ is the degree of f. The goal of this paper is to extend this result to the case when either $n = 7, 8$ or $n = 5, 6$ and $\mathrm{char}(K) > 3$.

2000 Mathematics Subject Classification: Primary 14H40; Secondary 14K05.

Key words and phrases. Hyperelliptic jacobians, Endomorphisms of abelian varieties, Supersingular abelian varieties.

1. Introduction

Let K be a field and K_a its algebraic closure. Assuming that $\mathrm{char}(K) = 0$, the author [25] proved that the jacobian $J(C) = J(C_f)$ of a hyperelliptic curve

$$C = C_f : y^2 = f(x)$$

has only trivial endomorphisms over K_a if the Galois group $\mathrm{Gal}(f)$ of the irreducible polynomial $f \in K[x]$ is "very big". Namely, if $n = \deg(f) \geq 5$ and $\mathrm{Gal}(f)$ is either the symmetric group $\mathbf{S}_n$ or the alternating group $\mathbf{A}_n$ then the ring $\mathrm{End}(J(C_f))$ of K_a-endomorphisms of $J(C_f)$ coincides with $\mathbf{Z}$. Later the author [25, 28] extended this result to the case of positive $\mathrm{char}(K) > 2$ but under the additional assumption that $n \geq 9$, i.e., the genus of C_f is greater or equal than 4. We refer the reader to [15, 16, 9, 10, 14, 11, 25, 26, 28, 29, 30] for a discussion of known results about, and examples of, hyperelliptic jacobians without complex multiplication.

The aim of the present paper is to extend this result to the case when either $n \geq 7$ or when $n \geq 5$ but $\mathrm{char}(K) > 3$. Notice that it is known [25] that in those cases either $\mathrm{End}(J(C)) = \mathbf{Z}$ or $J(C)$ is a supersingular abelian variety and the real problem is how to prove that $J(C)$ is *not* supersingular.

We also discuss the case of two-dimensional $J(C)$ in characteristic 3.

2. Main result

Throughout this paper we assume that K is a field of characteristic p different from 2. We fix its algebraic closure K_a and write $\mathrm{Gal}(K)$ for the absolute Galois group $\mathrm{Aut}(K_a/K)$.

Theorem 2.1. *Let K be a field with $p = \mathrm{char}(K) > 2$, K_a its algebraic closure, $f(x) \in K[x]$ an irreducible separable polynomial of degree n. Let us assume that $\mathrm{Gal}(f) = \mathbf{S}_n$ or $\mathbf{A}_n$. Suppose that n enjoys one of the following properties:*

(i) $n = 7$ *or* 8*;*
(ii) $n = 5$ *or* 6*. In addition,* $p = \mathrm{char}(K) > 3$.

Let C_f be the hyperelliptic curve $y^2 = f(x)$. Let $J(C_f)$ be its jacobian, $\mathrm{End}(J(C_f))$ *the ring of K_a-endomorphisms of $J(C_f)$. Then* $\mathrm{End}(J(C_f)) = \mathbf{Z}$.

Remark 2.2. Replacing K by a suitable finite separable extension, we may assume in the course of the proof of Theorem 2.1 that $\mathrm{Gal}(f) = \mathbf{A}_n$. Taking into account that $\mathbf{A}_n$ is simple non-abelian and replacing K by its abelian extension obtained by adjoining to K all 2-power roots of unity, we may also assume that K contains all 2-power roots of unity.

Remark 2.3. Let $f(x) \in K[x]$ be an irreducible separable polynomial of *even* degree $n = 2m \geq 6$ such that $\mathrm{Gal}(f) = \mathbf{S}_n$. Let $\alpha \in K_a$ be a root of f and $K_1 = K(\alpha)$ be the corresponding subfield of K_a. We have

$$f(x) = (x - \alpha)f_1(x)$$

with $f_1(x) \in K_1[x]$. Clearly, $f_1(x)$ is an irreducible separable polynomial over K_1 of degree $n - 1 = 2m - 1$, whose Galois group is SS_{n-1}. It is also clear that the polynomials

$$h(x) = f_1(x + \alpha), h_1(x) = x^{n-1}h(1/x) \in K_1[x]$$

are irreducible separable of degree $n - 1$ with the same Galois group SS_{n-1}.

The standard substitution

$$x_1 = 1/(x - \alpha), y_1 = y/(x - \alpha)^m$$

establishes a birational isomorphism between C_f and a hyperelliptic curve

$$C_{h_1} : y_1^2 = h_1(x_1).$$

In light of results of [27, 30] and Remarks 2.2 and 2.3, our Theorem 2.1 is an immediate corollary of the following auxiliary statement.

Theorem 2.4. *Let K be a field with $p = \mathrm{char}(K) > 2$, K_a its algebraic closure, $f(x) \in K[x]$ an irreducible separable polynomial of degree n. Let us assume that n and the Galois group* $\mathrm{Gal}(f)$ *of f enjoy one of the following properties:*

- (i) $n = 5$ *and* $\mathrm{Gal}(f) = \mathbf{A}_5$*;*
- (ii) $n = 7$ *and* $\mathrm{Gal}(f) = \mathbf{A}_7$. *In addition,* $p = \mathrm{char}(K) > 3$*;*

Let C be the hyperelliptic curve $y^2 = f(x)$ and let $J(C)$ be the jacobian of C. Then $J(C)$ is not a supersingular abelian variety.

We will prove Theorem 2.4 in Section 3.

Throughout the paper we write $\mathrm{End}^0(X)$ for the endomorphism algebra $\mathrm{End}(X) \otimes \mathbf{Q}$ of an abelian variety X over an algebraically closed field F_a. Recall [25] that the semisimple $\mathbf{Q}$-algebra $\mathrm{End}^0(X)$ has dimension $(2\dim(X))^2$ if and only if $p := \mathrm{char}(F_a) \neq 0$ and X is a supersingular abelian variety. We write $\mathbf{H}_p$ is the quaternion $\mathbf{Q}$-algebra unramified exactly at p and ∞. It is well-known that if X is a supersingular abelian variety in characteristic p then $\mathrm{End}^0(X)$ is isomorphic to the matrix algebra $\mathrm{M}_g(\mathbf{H}_p)$ of size $g := \dim(X)$ over $\mathbf{H}_p$. We will use freely these facts throughout the paper.

3. Proof of Theorem 2.4

We deduce Theorem 2.4 from the following statement.

Theorem 3.1. *Let K be a field with $p = \mathrm{char}(K) > 2$, K_a its algebraic closure, Let $n = q$ be an odd prime, $f(x) \in K[x]$ an irreducible separable polynomial of degree q. Let us assume that the Galois group* $\mathrm{Gal}(f)$ *of f is $\text{Ł}_2(q) := \mathrm{PSL}_2(\mathbf{F}_q)$, and that it acts doubly transitively on the roots of f. Suppose that either $q = 5$ or $q = 7$. Let C be the hyperelliptic curve $y^2 = f(x)$ and let $J(C)$ be the jacobian of C.*

If $J(C)$ is a supersingular abelian variety then $n = 5$ and $p = 3$.

Proof of Theorem 2.4 (modulo Theorem 3.1). If $n = 5$ then $\mathbf{A}_5 \cong \text{Ł}_2(5)$ and we are done.

Suppose that $n = 7$. It is well-known that the simple non-abelian group $\text{Ł}_2(7) \cong \text{Ł}_3(2) := \mathrm{PSL}_3(\mathbf{F}_2)$ acts doubly transitively on the 7-element projective plane $\mathbf{P}^2(\mathbf{F}_2)$ and therefore is isomorphic to a doubly transitive subgroup of $\mathbf{A}_7$. Hence there exists a finite algebraic extension K_1 of K such that the Galois group of f over K_1 is $\text{Ł}_2(7)$ acting doubly transitively on the roots of $f(x)$. Applying Theorem 3.1 to K_1 and f, we conclude that if $3 \ne \mathrm{char}(K_1) = \mathrm{char}(K) = p$ then $J(C)$ is not supersingular. $\square$

The following results will be used in order to prove Theorem 3.1.

Lemma 3.2. *Let K be a field with* $\mathrm{char}(K) \ne 2$ *K_a its algebraic closure,* $\mathrm{Gal}(K) = \mathrm{Aut}(K_a)$ *the Galois group of K. Let $f(x) \in K[x]$ be an irreducible separable polynomial of odd degree n. Let us assume that $n \ge 5$ and the Galois group* $\mathrm{Gal}(f)$ *of f acts doubly transitively on the roots of $f(x)$. Let C be the hyperelliptic curve $y^2 = f(x)$ and let $J(C)$ be the jacobian of C. Let $J(C)_2$ be the group of points of order 2 in $J(C)(K_a)$ viewed as $\mathbf{F}_2$-vector space provided with a natural structure of* $\mathrm{Gal}(K)$*-module. Then the image of* $\mathrm{Gal}(K)$ *in* $\mathrm{Aut}_{\mathbf{F}_2}(J(C)_2)$ *is isomorphic to* $\mathrm{Gal}(f)$ *and*

$$\mathrm{End}_{\mathrm{Gal}(K)}(J(C)_2) = \mathrm{End}_{\mathrm{Gal}(f)}(J(C)_2) = \mathbf{F}_2.$$

Theorem 3.3. *Let F be a field with characteristic $p > 2$ and assume that F contains all 2-power roots of unity. Let F_a be an algebraic closure of F.*

Let $G \ne \{1\}$ be a finite perfect group.

Suppose that g is a positive integer, X is a supersingular g-dimensional abelian variety defined over F. Let $\mathrm{End}(X)$ *is the ring of all F_a-endomorphisms of X and* $\mathrm{End}^0(X) = \mathrm{End}(X) \otimes \mathbf{Q}$.

Let us assume that the image of $\mathrm{Gal}(F)$ *in* $\mathrm{Aut}(X_2)$ *is isomorphic to G and the corresponding faithful representation*

$$\bar{\rho} : G \hookrightarrow \mathrm{Aut}(X_2) \cong \mathrm{GL}(2g, \mathbf{F}_2)$$

satisfies

$$\mathrm{End}_G X_2 = \mathbf{F}_2.$$

Then there exists a surjective group homomorphism

$$\pi_1 : G_1 \twoheadrightarrow G$$

enjoying the following properties:

(a) *The group G_1 is a perfect finite group. The kernel of π_1 is an elementary abelian 2-group.*

(b) *One may lift $\bar{\rho}\pi_1 : G_1 \to \mathrm{Aut}(X_2)$ to a faithful absolutely irreducible symplectic representation*

$$\rho : G_1 \hookrightarrow \mathrm{Aut}_{\mathbf{Q}_2}(V_2(X))$$

of G_1 *over* $\mathbf{Q}_2$ *in such a way that the following conditions hold:*

- *The character* χ *of* ρ *takes values in* $\mathbf{Q}$*;*
- $\rho(G_1) \subset (\mathrm{End}^0(X))^*$*;*
- *The homomorphism from the group algebra* $\mathbf{Q}[G_1]$ *to* $\mathrm{End}^0(X)$ *induced by* ρ *is surjective and identifies* $\mathrm{End}^0(X) \cong \mathrm{M}_g(\mathbf{H}_p)$ *with the direct summand of* $\mathbf{Q}[G_1]$ *attached to* χ*.*

(c) p *divides the order of* G *and* $p \le 2g+1$*.*

(d) *Suppose that either every homomorphism from* G *to* $\mathrm{GL}(g-1, \mathbf{F}_2)$ *is trivial or the* G*-module* X_2 *is very simple in the sense of* [27, 28, 31]*. Then* $\ker \pi_1$ *is a central cyclic group of order* 1 *or* 2*.*

Lemma 3.4. *Let* p *be an odd prime. Let* q *be an odd prime and* $\Gamma = \mathrm{SL}_2(\mathbf{F}_q)$ *or* $\mathrm{PSL}_2(\mathbf{F}_q)$*. Suppose that* $q = 5$ *or* 7 *and let us put* $g = \frac{q-1}{2}$*. Suppose that* $\mathbf{Q}[\Gamma]$ *contains a direct summand isomorphic to the matrix algebra* $\mathrm{M}_g(\mathbf{H}_p)$*. Then* $p = 3$ *and* $q = 5$*.*

Theorem 3.3 and Lemmas will be proven in Sections 5 and 4.

Proof of Theorem 3.1 (modulo Theorem 3.3 and Lemmas 3.2 and 3.4). Let us put

$$X = J(C), G = \mathrm{PSL}_2(\mathbf{F}_q), g = \frac{q-1}{2}.$$

Clearly, either $q = 5, g = 2$ or $q = 7, g = 3$. In both cases $g = \dim(X)$, the group G is simple and $\mathrm{GL}(g-1, \mathbf{F}_2)$ is solvable. It follows that every homomorphism from G to $\mathrm{GL}(g-1, \mathbf{F}_2)$ is trivial. It follows from Lemma 3.2 that the image of $\mathrm{Gal}(K)$ in $\mathrm{Aut}(X_2)$ is isomorphic to G and the corresponding faithful representation

$$\bar\rho : G \hookrightarrow \mathrm{Aut}(X_2) \cong \mathrm{GL}(2g, \mathbf{F}_2)$$

satisfies

$$\mathrm{End}_G X_2 = \mathbf{F}_2.$$

Let us assume that X is supersingular. We need to get a contradiction.

Applying Theorem 3.3, we conclude that there exist a finite perfect group G_1 and a surjective homomorphism

$$\pi_1 : G_1 \twoheadrightarrow G = \mathrm{PSL}_2(\mathbf{F}_q)$$

enjoying the following properties.

(i) Either $G_1 \cong G$ or $Z_1 = \ker(\pi_1)$ is a central subgroup of order 2 in G_1;

(ii) There exists a direct summand of $\mathbf{Q}[G_1]$ isomorphic to $\mathrm{M}_g(\mathbf{H}_p)$).

The well-known description of central extensions of $\mathrm{PSL}_2(\mathbf{F}_q)$ when q is an odd prime [4, §4.15, Prop. 4.233] implies that either $G_1 = \mathrm{PSL}_2(\mathbf{F}_q)$ or $G_1 = \mathrm{SL}_2(\mathbf{F}_q)$. Applying Lemma 3.4, we arrive to the desired contradiction. $\square$

4. PROOF OF LEMMAS 3.2 AND 3.4

We start with some auxiliary constructions related to the permutation groups [12, 17, 7].

Let B be a finite set consisting of $n \ge 5$ elements. We write $\mathrm{Perm}(B)$ for the group of permutations of B. A choice of ordering on B gives rise to an isomorphism $\mathrm{Perm}(B) \cong \mathbf{S}_n$. Let us assume that n is *odd* and consider the permutation module $\mathbf{F}_2^B$: the $\mathbf{F}_2$-vector space of all functions $\varphi : B \to \mathbf{F}_2$. The space $\mathbf{F}_2^B$ carries a natural structure of $\mathrm{Perm}(B)$-module and contains the stable hyperplane $Q_B := (\mathbf{F}_2^B)^0$ of functions φ with $\sum_{\alpha\in B}\varphi(\alpha)=0$. Clearly, Q_B carries a natural structure of faithful $\mathrm{Perm}(B)$-module. For each permutation group $H\subset \mathrm{Perm}(B)$ the corresponding H-module is called the *heart* of the permutation representation of H on B over $\mathbf{F}_2$ [12, 17, 7].

Lemma 4.1. $\mathrm{End}_H(Q_B)=\mathbf{F}_2$ *if n is odd and H acts 2-transitively on B.*

Proof. See Satz 4 in [12]. $\square$

Proof of Lemma 3.2. Suppose $f(x)\in K[x]$ is a polynomial of odd degree $n\ge 5$ without multiple roots and $X:=J(C_f)$ is the jacobian of $C=C_f: y^2=f(x)$. It is well-known that $g:=\dim(X)=\frac{n-1}{2}$. It is also well-known (see for instance Sect. 5 of [27]) that the image of $\mathrm{Gal}(K)\to \mathrm{Aut}(X_2)$ is isomorphic to $\mathrm{Gal}(f)$. More precisely, let $\mathfrak{R}\subset K_a$ be the n-element set of roots of f, let $K(\mathfrak{R})$ be the splitting field of f and $\mathrm{Gal}(f)=\mathrm{Gal}(K(\mathfrak{R})/K)$ the Galois group of f, viewed as a subgroup of of the group $\mathrm{Perm}(\mathfrak{R})$ of all permutations of $\mathfrak{R}$. We have $\mathrm{Gal}(f)\subset \mathrm{Perm}(\mathfrak{R})$. It is well-known (see for instance, Th. 5.1 on p. 478 of [27]) that $\mathrm{Gal}(K)\to \mathrm{Aut}(X_2)$ factors through the canonical surjection $\mathrm{Gal}(K)\twoheadrightarrow \mathrm{Gal}(K(\mathfrak{R})/K)=\mathrm{Gal}(f)$ and the $\mathrm{Gal}(f)$-modules X_2 and $Q_{\mathfrak{R}}$ are isomorphic. In particular,

$$\mathrm{End}_{\mathrm{Gal}(K)}(X_2)=\mathrm{End}_{\mathrm{Gal}(f)}(X_2)=\mathrm{End}_{\mathrm{Gal}(f)}(Q_{\mathfrak{R}}).$$

Assuming that $\mathrm{Gal}(f)$ acts doubly transitively on $\mathfrak{R}$ and applying Lemma 4.1, we conclude that

$$\mathrm{End}_{\mathrm{Gal}(f)}(X_2)=\mathrm{End}_{\mathrm{Gal}(f)}(Q_{\mathfrak{R}})=\mathbf{F}_2.$$

$\square$

Remark 4.2. The assertion of Lemma 3.2 is implicitly contained in the proof of Prop. 3 in [16].

Proof of Lemma 3.4. It is known [8, corollary on p. 4] that $\mathbf{Q}[\mathrm{PSL}_2(\mathbf{F}_q)]$ is a direct product of matrix algebras (for all power primes q). Since $\ker(\mathrm{SL}_2(\mathbf{F}_q)\twoheadrightarrow \mathrm{PSL}_2(\mathbf{F}_q))$ is the only proper normal subgroup in $\mathrm{SL}_2(\mathbf{F}_q)$, it suffices to deal only with the group $\mathrm{SL}_2(\mathbf{F}_q)$ with $q=5, g=2$ or $q=7, g=3$ and consider only direct summands of $\mathbf{Q}[\mathrm{SL}_2(\mathbf{F}_q)]$ that correspond (in the sense of Lemma 24.7 on p. 124 of [2]) to *faithful* irreducible characters of degree $q-1$ with values in $\mathbf{Q}$.

Let χ be an *irreducible faithful irreducible* character of degree $q-1$ with values in $\mathbf{Q}$. Then (in the notations of [2, §38]) $\chi=\theta_j$ where j is an integer with $1\le j\le \frac{q-1}{2}$. If z is the only nontrivial central element of $\mathrm{SL}_2(\mathbf{F}_q)$ then $\theta_j(z)=(-1)^j(q-1)$. The faithfulness of χ implies (thanks to Lemma 2.19 [6]) that $\theta_j(z)\ne q-1$, i.e. j is odd. Let $b\in \mathrm{SL}_2(\mathbf{F}_q)$ be an element of order q and σ a primitive $q+1$th root of unity. Then [2, p. 228]

$$\chi(b)=\theta_j(b)=-(\sigma^j+\sigma^{-j}).$$

Assume that $q=7$. Then either $j=1$ or $j=3$. Also $q+1=8$ and we may choose

$$\sigma=\frac{1+\sqrt{-1}}{\sqrt{2}}.$$

Then if $j=1$ then $\chi(b)=-\sqrt{2}$ and if $j=3$ then $\chi(b)=\sqrt{2}$. In both cases $\chi(b)$ does not lie in $\mathbf{Q}$. It follows that $\mathbf{Q}[\mathrm{SL}_2(\mathbf{F}_7)]$ does not have direct summands isomorphic to the matrix algebras of size 3 over quaternion $\mathbf{Q}$-algebras (including $\mathbf{H}_p$).

Assume that $q = 5$. Then $j = 1$ and $\chi = \theta_1$. Then $q+1 = 6$ and the multiplicative order n of σ^j equals $6 = 2 \cdot 3$. Also $\sigma^{2j} = \sigma^2$ is a primitive cubic root of unity. Let D be the direct summand of $\mathbf{Q}[\mathrm{SL}_2(\mathbf{F}_5)]$ attached to χ. It follows from the case (c) of Theorem on p. 4 of [8] (see also [3, Th. 6.1(ii)] (with $\epsilon = \delta = 1$)) that D is isomorphic to to the matrix algebra $\mathrm{M}_2(\mathbf{H})$ where H is a quaternion $\mathbf{Q}$-algebra ramified (exactly) at ∞ and 3. (This means that $H \cong \mathbf{H}_3$ and $D \cong \mathrm{M}_2(\mathbf{H}_3)$.) It follows that if D is isomorphic to $\mathrm{M}_2(\mathbf{H}_p)$ then $p = 3$. $\square$

5. NOT SUPERSINGULARITY

We keep all the notations and assumptions of Theorem 3.3.

We write $T_2(X)$ for the 2-adic Tate module of X and

$$\rho_{2,X} : \mathrm{Gal}(F) \to \mathrm{Aut}_{\mathbf{Z}_2}(T_2(X))$$

for the corresponding 2-adic representation. It is well-known that $T_2(X)$ is a free $\mathbf{Z}_2$-module of rank $2\mathrm{dim}(X) = 2g$ and

$$X_2 = T_2(X)/2T_2(X)$$

(the equality of Galois modules). Let us put

$$H = \rho_{2,X}(\mathrm{Gal}(F)) \subset \mathrm{Aut}_{\mathbf{Z}_2}(T_2(X)).$$

Clearly, the natural homomorphism

$$\bar{\rho}_{2,X} : \mathrm{Gal}(F) \to \mathrm{Aut}(X_2)$$

defining the Galois action on the points of order 2 is the composition of $\rho_{2,X}$ and (surjective) reduction map modulo 2

$$\mathrm{Aut}_{\mathbf{Z}_2}(T_2(X)) \to \mathrm{Aut}(X_2).$$

This gives us a natural (continuous) *surjection*

$$\pi : H \to \bar{\rho}_{2,X}(\mathrm{Gal}(F)) \cong G,$$

whose kernel consists of elements of $1 + 2\mathrm{End}_{\mathbf{Z}_2}(T_2(X))$. The choice of polarization on X gives rise to a non-degenerate alternating bilinear form (Riemann form) [18]

$$e : V_2(X) \times V_2(X) \to \mathbf{Q}_2(1) \cong \mathbf{Q}_2.$$

Since F contains all 2-power roots of unity, e is $\mathrm{Gal}(F)$-invariant and therefore is H-invariant. In particular,

$$H \subset \mathrm{Sp}(V_2(X), e) \subset \mathrm{SL}(V_2(X)).$$

Here $\mathrm{Sp}(V_2(X), e)$ is the symplectic group attached to e. In particular, the H-module $V_2(X)$ is symplectic.

There exists a finite Galois extension L of F such that all endomorphisms of X are defined over L. Clearly, $\mathrm{Gal}(L) = \mathrm{Gal}(F_a/L)$ is an open normal subgroup of finite index in $\mathrm{Gal}(F)$ and

$$H' = \rho_{2,X}(\mathrm{Gal}(L)) \subset \mathrm{Aut}_{\mathbf{Z}_2}(T_2(X)) \subset \mathrm{Aut}_{\mathbf{Q}_2}(V_2(X)))$$

is an open normal subgroup of finite index in H. We write $\mathrm{End}^0(X)$ for the $\mathbf{Q}$-algebra $\mathrm{End}(X) \otimes \mathbf{Q}$ of endomorphisms of X.

There exists a finite Galois extension L of F such that all endomorphisms of X are defined over L. We write $\mathrm{End}^0(X)$ for the $\mathbf{Q}$-algebra $\mathrm{End}(X) \otimes \mathbf{Q}$ of endomorphisms of X.

Since X is supersingular,

$$\dim_{\mathbf{Q}}\mathrm{End}^0(X) = (2\dim(X))^2 = (2g)^2.$$

Recall ([18]) that the natural map

$$\mathrm{End}^0(X)\otimes_{\mathbf{Q}}\mathbf{Q}_2 \to \mathrm{End}_{\mathbf{Q}_2}V_2(X)$$

is an embedding. Dimension arguments imply that

$$\mathrm{End}^0(X)\otimes_{\mathbf{Q}}\mathbf{Q}_2 = \mathrm{End}_{\mathbf{Q}_2}V_2(X).$$

Since all endomorphisms of X are defined over L, the image

$$\rho_{2,X}(\mathrm{Gal}(L)) \subset \rho_{2,X}(\mathrm{Gal}(F)) \subset \mathrm{Aut}_{\mathbf{Z}_2}(T_2(X)) \subset \mathrm{Aut}_{\mathbf{Q}_2}(V_2(X))$$

commutes with $\mathrm{End}^0(X)$. This implies that $\rho_{2,X}(\mathrm{Gal}(L))$ commutes with $\mathrm{End}_{\mathbf{Q}_2}V_2(X)$ and therefore consists of scalars. Since

$$\rho_{2,X}(\mathrm{Gal}(L)) \subset \rho_{2,X}(\mathrm{Gal}(F)) \subset \mathrm{SL}(V_2(X)),$$

$\rho_{2,X}(\mathrm{Gal}(L))$ is a finite group. Since $\mathrm{Gal}(L)$ is a subgroup of finite index in $\mathrm{Gal}(F)$, the group $H = \rho_{2,X}(\mathrm{Gal}(F))$ is also finite. In particular, the kernel of the reduction map modulo 2

$$\mathrm{Aut}_{\mathbf{Z}_2}T_2(X) \supset H \to G \subset \mathrm{Aut}(X_2)$$

consists of periodic elements and, thanks to Minkowski-Serre Lemma [23], $Z := \ker(\pi : H \to G)$ has exponent 1 or 2. In particular, Z is commutative. Since

$$Z \subset H \subset \mathrm{Sp}(V_2(X)) \cong \mathrm{Sp}(2g, \mathbf{Q}_2),$$

Z is a $\mathbf{F}_2$-vector space of dimension $\le g$.

Let G_1 be a minimal subgroup of H such that $\pi(G_1) = G$. (Since H is finite, such G_1 always exists.) Since G is perfect, G_1 is also perfect. (Otherwise, we may replace G_1 by smaller $[G_1, G_1]$.) Clearly,

$$Z_1 := \ker(\pi : G_1 \twoheadrightarrow G) \subset Z$$

is also a $\mathbf{F}_2$-vector space of dimension $\le g$. We have

$$Z_1 \subset G_1 \subset H \subset \mathrm{Sp}(V_2(X)) \cong \mathrm{Sp}(2g, \mathbf{Q}_2).$$

In particular, the symplectic G_1-module is a lifting of the $G_1(\twoheadrightarrow G)$-module X_2.

I claim that the natural representation of G_1 in the $2g$-dimensional $\mathbf{Q}_2$-vector space $V_2(X)$ is absolutely irreducible. Indeed, let us put

$$E := \mathrm{End}_{G_1}(V_2(X)) \subset \mathrm{End}_{\mathbf{Q}_2}(V_2(X)).$$

Clearly,

$$O_E = E \bigcap \mathrm{End}_{\mathbf{Z}_2}(T_2(X)) \subset \mathrm{End}_{\mathbf{Z}_2}(T_2(X))$$

is a $\mathbf{Z}_2$-algebra that is a free $\mathbf{Z}_2$-module, whose $\mathbf{Z}_2$-rank coincides with $\dim_{\mathbf{Q}_2}(E)$. Notice that O_E is a *pure* $\mathbf{Z}_2$-submodule in $\mathrm{End}_{\mathbf{Z}_2}(T_2(X))$, i.e. the quotient $\mathrm{End}_{\mathbf{Z}_2}(T_2(X))/O_E$ is a torsion-free (finitely generated) $\mathbf{Z}_2$-module and therefore a free $\mathbf{Z}_2$-module of finite rank. It follows that the natural map

$$O_E/2O_E \to \mathrm{End}_{\mathbf{Z}_2}(T_2(X))/2\mathrm{End}_{\mathbf{Z}_2}(T_2(X)) = \mathrm{End}_{\mathbf{F}_2}(X_2)$$

is an embedding. Clearly, the image of $O_E/2O_E$ in $\mathrm{End}_{\mathbf{F}_2}(X_2)$ lies in $\mathrm{End}_G(X_2)$. Since $\mathrm{End}_G(X_2) = \mathbf{F}_2$, we conclude that the rank of the free $\mathbf{Z}_2$-module O_E is 1, i.e. $\dim_{\mathbf{Q}_2}(E) = 1$. This means that $E = \mathbf{Q}_2$, i.e. the G_1-module $V_2(X)$ is absolutely simple.

Let $\chi : G_1 \to \mathbf{Q}_2$ be the character of the absolutely irreducible faithful representation of G_1 in $V_2(X)$. Clearly, χ is a faithful (absolutely) irreducible character of degree $2g$. We need to prove that $\chi(G_1) \subset \mathbf{Q}$.

Let $F_1 \subset F_a$ be the subfield of invariants of the subgroup

$$\{\sigma \in \mathrm{Gal}(F) \mid \rho_{2,X}(\sigma) \in G_1\} \subset \mathrm{Gal}(F).$$

Clearly, F_1 is a finite separable algebraic extension of F and

$$G_1 = \rho_{2,X}(\mathrm{Gal}(F_1)).$$

Clearly, the image

$$\bar{\rho}_{2,X}(\mathrm{Gal}(F_1)) \subset \mathrm{Aut}(X_2)$$

coincides with

$$\pi\rho_{2,X}(\mathrm{Gal}(F_1)) = \pi(G_1) = \pi_1(G_1) = G \subset \mathrm{Aut}(X_2).$$

Let L_1 be the finite Galois extension of F_1 attached to

$$\rho_{2,X} : \mathrm{Gal}(F_1)) \to \mathrm{Aut}(T_2(X)).$$

Clearly, $\mathrm{Gal}(L_1/F_1) = G_1$. In addition, all 2-power torsion points of X are defined over L_1. It follows (see [22]) that all the endomorphisms of X are defined over L_1. On the other hand, I claim that the ring $\mathrm{End}_{F_1}(X)$ of F_1-endomorphisms of X coincides with $\mathbf{Z}$. Indeed, there is a natural embedding

$$\mathrm{End}_{F_1}(X) \otimes \mathbf{Z}/2\mathbf{Z} \hookrightarrow \mathrm{End}_{\mathrm{Gal}(F_1)}(X_2) = \mathbf{F}_2$$

that implies that the rank of the free $\mathbf{Z}$-module $\mathrm{End}_{F_1}(X)$ does not exceed 1 and therefore equals 1, i.e. $\mathrm{End}_{F_1}(X) = \mathbf{Z}$.

Since all the endomorphisms of X are defined over L_1, there is a natural homomorphism

$$\kappa : G_1 = \mathrm{Gal}(L_1/F_1) \to \mathrm{Aut}(\mathrm{End}(X))$$

such that

$$\mathrm{End}_{F_1}(X) = \{u \in \mathrm{End}(X) \mid \kappa(\sigma)u = u \quad \forall \sigma \in \mathrm{Gal}(L_1/F_1) = G_1\}$$

and

$$\sigma(ux) = (\kappa(\sigma)u)(\sigma(x)) \quad \forall x \in X(L_1), u \in \mathrm{End}(X), \sigma \in \mathrm{Gal}(L_1/F_1) = G_1.$$

Further we write ${}^{\kappa(\sigma)}u$ for $\kappa(\sigma)(u)$. Since $\mathrm{End}_{F_1}(X) = \mathbf{Z}$, we conclude that

$$\mathbf{Z} = \{u \in \mathrm{End}(X) \mid {}^{\kappa(\sigma)}u = u \quad \forall \sigma \in \mathrm{Gal}(L_1/F_1) = G_1\}.$$

Since all 2-power torsion points of X defined over L_1,

$$\sigma(ux) = {}^{\kappa(\sigma)}u(\sigma(x)) \quad \forall x \in T_2(X), u \in \mathrm{End}(X), \sigma \in G_1.$$

Since $\mathrm{Aut}(\mathrm{End}(X)) \subset \mathrm{Aut}(\mathrm{End}^0(X))$, one may view κ as

$$\kappa : G_1 = \mathrm{Gal}(L_1/F_1) \to \mathrm{Aut}(\mathrm{End}^0(X)), \quad u \mapsto {}^{\kappa(\sigma)}u, \quad u \in \mathrm{End}^0(X), \sigma \in G_1$$

and we have

$$\mathbf{Q} = \{u \in \mathrm{End}^0(X) \mid {}^{\kappa(\sigma)}u = u \quad \forall \sigma \in \mathrm{Gal}(L_1/F_1) = G_1\}$$

and

$$\sigma(ux) = {}^{k(\sigma)}u(\sigma(x)) \quad \forall x \in V_2(X), u \in \mathrm{End}^0(X), \sigma \in G_1.$$

Recall that

$$\mathrm{End}^0(X) \subset \mathrm{End}^0(X) \otimes_{\mathbf{Q}} \mathbf{Q}_2 = \mathrm{End}_{\mathbf{Q}_2}(V_2(X))$$

and

$$G_1 \subset \mathrm{GL}(V_2(X)) = (\mathrm{End}_{\mathbf{Q}_2}(V_2(X)))^*.$$

It follows that

$$\sigma u \sigma^{-1} = {}^{k(\sigma)}u \quad \forall u \in \mathrm{End}^0(X), \sigma \in G_1.$$

By Skolem-Noether theorem, every automorphism of the central simple $\mathbf{Q}$-algebra $\mathrm{End}^0(X) \cong \mathrm{M}_g(\mathbf{H}_p)$ is an inner one. This implies that for each $\sigma \in G_1$ there exists $w_\sigma \in \mathrm{End}^0(X)^*$ such that

$$\sigma u \sigma^{-1} = {}^{k(\sigma)}u = w_\sigma u w_\sigma^{-1} \quad \forall u \in \mathrm{End}^0(X).$$

Since the center of $\mathrm{End}^0(X)$ is $\mathbf{Q}$, the choice of w_σ is unique up to multiplication by a non-zero rational number. This implies that $w_\sigma w_\tau$ equals $w_{\sigma\tau}$ times a non-zero rational number.

Let us put

$$c'_\sigma = \sigma w_\sigma^{-1} \in (\mathrm{End}_{\mathbf{Q}_2}(V_2(X)))^*.$$

Clearly, each c'_σ commutes with $\mathrm{End}^0(X)$ and therefore with $\mathrm{End}^0(X) \otimes_{\mathbf{Q}} \mathbf{Q}_2 = \mathrm{End}_{\mathbf{Q}_2}(V_2(X))$. It follows that all c'_σ are scalars, i.e. lie in $\mathbf{Q}_2^*\mathrm{Id}$. (Here Id is the identity map on $V_2(X)$.) Clearly, the image

$$c_\sigma \in \mathbf{Q}_2^*\mathrm{Id}/\mathbf{Q}^*\mathrm{Id} \cong {\mathbf{Q}_2}^*/\mathbf{Q}^*$$

of c'_σ in ${\mathbf{Q}_2}^*/\mathbf{Q}^*$ does not depend on the choice of w_σ. It is also clear that the map

$$G_1 \to {\mathbf{Q}_2}^*/\mathbf{Q}^*, \sigma \mapsto c'_\sigma$$

is a group homomorphism. Since G_1 is perfect and ${\mathbf{Q}_2}^*/\mathbf{Q}^*$ is commutative, this homomorphism is trivial, i.e. $c_\sigma = 1$ for all $\sigma \in G_1$. This means that

$$c_\sigma \in \mathbf{Q}^*\mathrm{Id} \quad \forall \sigma \in G_1$$

and therefore

$$\sigma = (c'_\sigma)^{-1} w_\sigma \in \mathrm{End}^0(X)^* \quad \forall \sigma \in G_1.$$

Recall [18] that if one view an element $u \in \mathrm{End}^0(X)$ as linear operator in $V_2(X)$ then the characteristic polynomial $P_u(t)$ of u has rational coefficients; in particular, the trace of u is a rational number. It follows that $\chi(G_1) \subset \mathbf{Q}$.

Let M be the image of

$$\mathbf{Q}[G_1] \to \mathrm{End}^0(X).$$

Clearly, $M \otimes_{\mathbf{Q}} \mathbf{Q}_2$ coincides with the image of

$$\mathbf{Q}_2[G_1] \to \mathrm{End}^0(X) \otimes_{\mathbf{Q}} \mathbf{Q}_2 = \mathrm{End}_{\mathbf{Q}_2}(V_2(X)).$$

Since the G_1-module $V_2(X)$ is absolutely simple,

$$\mathbf{Q}_2[G_1] \to \mathrm{End}_{\mathbf{Q}_2}(V_2(X))$$

is surjective. This implies that

$$\dim_{\mathbf{Q}}(M) = \dim_{\mathbf{Q}}(\mathrm{End}^0(X))$$

and therefore, $M = \mathrm{End}^0(X)$, i.e. $\mathbf{Q}[G_1] \to \mathrm{End}^0(X)$ is surjective. The semisimplicity of $\mathbf{Q}[G_1]$ allows us to identify $\mathrm{End}^0(X)$ with a direct summand of $\mathbf{Q}[G_1]$.

If ℓ is a prime number that does not divide order of G_1 then it is well-known that the group algebra $\mathbf{Q}_\ell[G_1]$ is a direct product of matrix algebras over (commutative) fields. It follows that p divides order of G_1. Since $\#(G_1)$ equals $\#G$ times a power of 2 and p is odd, we conclude that p divides $\#G$. In particular, G_1 contains an element u of exact order p. Since

$$u \in G_1 \subset \mathrm{End}^0(X) \subset \mathrm{End}_{\mathbf{Q}_2}(V_2(X)),$$

$P_u(t)$ is a polynomial of degree $2g$ with rational coefficients and one of its roots is a primitive pth root of unity. It follows that $P_u(t)$ is divisible in $\mathbf{Q}[t]$ by the pth cyclotomic polynomial $\Phi_p(t) = \frac{t^p-1}{t-1}$. Since the degree of Φ_p is $p-1$, we conclude that the degree $2g$ of $P_u(t)$ is greater or equal than $p-1$, i.e. $2g \ge p-1$.

Assume for a while that the G-module X_2 is very simple. Since $G_1 \to G$ is surjective, the G_1-module X_2 and its lifting $V_2(X)$ are also very simple G_1-modules [28, Remark 5.2 (i,v(a))]. Since Z_1 is normal in G_1, we conclude, thanks to [28, Remark 5.2(vii)] that either the Z_1-module $V_2(X)$ is absolutely simple or Z_1 consists of scalars. Since Z_1 is a finite commutative group, it does not admit absolutely irreducible representations of dimension > 1. Since $\dim_{\mathbf{Q}_2}(V_2(X)) = 2g > 1$, we conclude that Z_1 consists scalars; in particular, Z_1 is a central subgroup in G_1. Since

$$Z_1 \subset G_1 \subset \mathrm{Sp}(V_2(X)) \cong \mathrm{Sp}(2g, \mathbf{Q}_2),$$

either $Z = \{1\}$ or $Z = \{\pm 1\}$. This implies that Z_1 is a cyclic group of order 1 or 2.

Further we no longer assume that the G-module X_2 is very simple. Assume instead that every homomorphism from Z to $\mathrm{GL}(g-1, \mathbf{F}_2)$ is trivial. I claim that in this case Z is again a central subgroup of G_1. Indeed, the short exact sequence

$$1 \to Z \hookrightarrow G_1 \twoheadrightarrow G \to 1$$

defines, in light of commutativeness of Z, a natural homomorphism

$$\eta : G \to \mathrm{Aut}(Z)$$

which is trivial if and only if Z is central in G_1. Clearly, $\eta(G)$ is a finite perfect group. Recall that Z is an elementary 2-group, i.e. $Z \cong \mathbf{F}_2^r$ for some nonnegative integer r. Clearly, we may assume that $r \ge 1$ and therefore $\mathrm{Aut}(Z) \cong \mathrm{GL}(r, \mathbf{F}_2)$. If $r \le g-1$ then we are done. Suppose that $r = g$. Then Z must contain

$$\{\pm 1\} \subset \mathrm{Sp}(V_2(X).$$

Since $\{\pm 1\}$ is a central subgroup of G_1, the elements of $\eta(G) \subset \mathrm{Aut}(Z)$ act trivially on $\{\pm 1\}$. Since the quotient $Z/\{\pm 1\}$ has $\mathbf{F}_2$-dimension $g-1$, elements of $\eta(G)$ act trivially on $Z/\{\pm 1\}$. This implies that $\eta(G)$ is isomorphic to a subgroup of the commutative group $\mathrm{Hom}(Z/\{\pm 1\}, \{\pm 1\}$. Since $\eta(G)$ is perfect, we conclude that $\eta(G) = \{1\}$, i.e. Z is a central subgroup and therefore is either $\{1\}$ or $\{\pm 1\}$.

6. Hyperelliptic two-dimensional jacobians in characteristic 3

Throughout this section K is a field of characteristic $p = 3$ and K_a its algebraic closure, $n = 5$ or 6,

$$f(x) = \sum_{i=0}^{n} a_i x^i \in K[x]$$

a separable polynomial of degree n, i.e. all $a_i \in K, a_n \ne 0$ and f has no multiple roots. We write $\mathrm{Gal}(f) \subset SS_n$ for the Galois group of f over K.

Let C_f be the hyperelliptic curve $y^2 = f(x)$ over K_a.

Lemma 6.1. *Suppose that* $n = \deg(f) = 5$ *and* $a_4 = 0$.

(i) *The jacobian $J(C_f)$ of C_f is a supersingular abelian variety over K_a if and only if $a_1 = a_2 = 0$, i.e.*
$$f(x) = a_5x^5 + a_3x^3 + a_0.$$
If this is the case then $J(C_f)$ is isogenous but not isomorphic to a self-product of a supersingular elliptic curve.

(ii) *Suppose that $a_0 \neq 0$ (e.g., $f(x)$ is irreducible over K) and $J(C_f)$ is a supersingular abelian variety. Then* $\mathrm{Gal}(f) \subset \mathbf{A}_5$ *if and only if -1 is a square in K, i.e. K contains $\mathbf{F}_9$.*

Proof. Since $p = 3$, $f(x)^{(p-1)/2} = f(x)$. Let us consider the matrices
$$M := \begin{pmatrix} a_{p-1} & a_{p-2} \\ a_{2p-1} & a_{2p-2} \end{pmatrix} = \begin{pmatrix} a_2 & a_1 \\ a_5 & 0 \end{pmatrix}, M^{(3)} := \begin{pmatrix} a_2^3 & a_1^3 \\ a_5^3 & 0 \end{pmatrix}.$$
Extracting cubic roots from all entries of M one gets the Hasse–Witt/Cartier–Manin matrix $M^{(3)}$ of C (with respect to the standard basis in the space of differentials of the first kind) [13], [24], [5, p. 129]. Recall ([13, p. 78], [19], [24, Th. 3.1], [5, Lemma 1.1]) that the jacobian $J(C)$ is a supersingular abelian surface not isomorphic to a product of two supersingular elliptic curves if and only if $M \neq 0$ but
$$M^{(3)}M = 0.$$
Clearly, $M \neq 0$, because $a_5 \neq 0$. It is also clear that
$$\det(M^{(3)}M) = \det(M^{(3)})\det(M) = (-a_1^3a_5^3)(-a_1a_5) = a_1^4a_5^4.$$
Hence, if $M^{(3)}M = 0$ then $a_1 = 0$.

Suppose that $a_1 = 0$. Then
$$M = \begin{pmatrix} a_2 & 0 \\ a_5 & 0 \end{pmatrix}, M^{(3)} = \begin{pmatrix} a_2^3 & 0 \\ a_5^3 & 0 \end{pmatrix}, M^{(3)}M = \begin{pmatrix} a_2^4 & 0 \\ a_5^3a_2 & 0 \end{pmatrix}.$$
We conclude that $M^{(3)}M = 0$ if and only if $a_1 = a_2 = 0$. It follows that $J(C)$ is a supersingular abelian surface if and only if $a_1 = a_2 = 0$. Since $M \neq 0$, the jacobian $J(C)$ is not isomorphic to a product of two supersingular elliptic curves. This proves (i).

In order to prove (ii), let us assume that $J(C_f)$ is supersingular, i.e.,
$$f(x) = a_5X^5 + a_3x^3 + a_0.$$
We know that $a_0 \neq 0, a_5 \neq 0$. Let us put
$$h(x) := a_5^{-1}f(x) = x^5 + b_3x^3 + b_0$$
where $b_3 = a_3/a_5, b_0 = a_0/a_5$. Clearly, $b_0 \neq 0$ and the Galois groups of $f(x)$ and $h(x)$ coincide. So, it suffices to check that $\mathrm{Gal}(h) \subset \mathbf{A}_5$ if and only if -1 is a square in K.

The derivative $h'(x)$ of $h(x)$ is $5x^4 = -x^4$. Let $\alpha_1, \cdots, \alpha_5$ be the roots of h. Clearly,
$$\prod_{i=1}^{5} \alpha_i = -b_0.$$
It is well-known that the Galois group of h lies in the alternating group if and only if its discriminant
$$D = \prod_{i<j}(\alpha_i - \alpha_j)^2$$

is a square in K. On the other hand, it is also well-known that

$$\prod_{i=1}^{5} h'(\alpha_i) =: R(h,h') = (-1)^{\deg(h)(\deg(h)-1)/2} D.$$

(Here $R(h,h')$ is the resultant of h and h'.) It follows that

$$R(h,h') = \prod_{i=1}^{5}(-\alpha_i^4) = -(\prod_{i=1}^{5}\alpha_i)^4 = -(-b_0)^4 = -b_0^4$$

and therefore $D = -b_0^4$. Clearly, D is a square in K if and only if -1 is a square in K. □

Example 6.2 (Counterexamples for $\mathbf{A}_5$ and SS_5)**.** Let k be an algebraically closed field of characteristic $p = 3$. Let $K = k(z)$ be the field of rational functions in variable z with constant field k. We write $\overline{k(z)}$ for an algebraic closure of $k(z)$. According to Abhyankar [1], the Galois group of the polynomial

$$h(x) = x^5 - zx^2 + 1 \in k(z)[x] = K[x]$$

is $\mathbf{A}_5$ (see also [21, §3.3]). It follows that the Galois group of the polynomial

$$f(x) = x^5 h\left(\frac{1}{x}\right) = x^5 - zx^3 + 1 = \sum_{i=1}^{5} a_i x^i$$

is also $\mathbf{A}_5$. (Here $a_5 = 1, a_4 = a_2 = a_1 = 0, a_3 = -z, a_0 = 1$.)

Let us consider the hyperelliptic curve

$$C : y^2 = x^5 - zx^3 + 1$$

of genus 2 over $\overline{k(z)}$. It follows from Lemma 6.1 that the jacobian $J(C)$ of C is a supersingular abelian surface that is *not* isomorphic to a product of two supersingular elliptic curves. Hence $\mathrm{End}(J(C))$ is isomorphic to a certain order in the matrix algebra of size 2 over the quaternion $\mathbf{Q}$-algebra ramified exactly at 3 and ∞. See [5, proposition 2.19]) for an explicit description of this order.

Assume now that k is an algebraic closure of $\mathbf{F}_3$. Let us put

$$K_0 = \mathbf{F}_3(z) \subset K = k(z) \subset \overline{k(z)}.$$

Clearly, -1 is *not* a square in K_0 and $\overline{k(z)}$ is an algebraic closure of K_0. Also, $f(x) \in K_0[x]$. An elementary calculation (as in the proof of Lemma 6.1(ii)) shows that the discriminant of $f(x)$ is -1. This implies that the Galois group of $f(x)$ over K_0 does not lie in $\mathbf{A}_5$. It follows that the Galois group of $f(x) = x^5 - zx^3 + 1$ over K_0 is SS_5. However, as we have already seen, the jacobian of $y^2 = x^5 - zx^3 + 1$ is supersingular.

Theorem 6.3. *Let K be a field with* $\mathrm{char}(K) = 3$*, K_a its algebraic closure, $f(x) \in K[x]$ an irreducible separable polynomial of degree $n = 5$ or 6. Let us assume that the Galois group* $\mathrm{Gal}(f)$ *of f is the full symmetric group $\mathbf{S}_n$. Assume, in addition, that -1 is a square in K, i.e. K contains $\mathbf{F}_9$.*

Let $C = C_f$ be the hyperelliptic curve $y^2 = f(x)$. Let $J(C_f)$ be its jacobian, $\mathrm{End}(J(C_f))$ *the ring of K_a-endomorphisms of $J(C_f)$. Then* $\mathrm{End}(J(C_f)) = \mathbf{Z}$.

Proof of Theorem 6.3. Thanks to Remark 2.3, we may and will assume that $n = 5$. We have

$$f(x) = \sum_{i=0}^{5} a_i x^i \in K[x]$$

where all the coefficients $a_i \in K$ and $a_0 \neq 0$. Let us put

$$\gamma := \frac{a_4}{5a_0}, \quad h(x) := f(x - \gamma).$$

Clearly, $h(x) \in K[x]$ is an irreducible polynomial of degree 5 and $\mathrm{Gal}(h) = \mathrm{Gal}(f) = SS_5$. It is also clear that if

$$h(x) = \sum_{i=0}^{5} b_i x^i \in K[x]$$

then $b_4 = 0,\ b_5 = a_5 \neq 0$. The substitution $x_1 = x + \gamma,\ y_1 = y$ establishes a K-birational isomorphism between hyperelliptic curves $C = C_f : y^2 = f(x)$ and $C_1 = C_h : y_1^2 = h(x_1)$ and induces an isomorphism of the jacobians $J(C_f)$ and $J(C_h)$.

Suppose that $\mathrm{End}(J(C_f)) \neq \mathbf{Z}$. Then it follows from Theorem 2.1 of [25] that $J(C_f)$ is a supersingular abelian variety. It follows that $J(C_h) \cong J(C_f)$ is also a supersingular abelian variety. Applying Lemma 6.1(ii) to h, we conclude that $\mathrm{Gal}(h) \subset \mathbf{A}_5$, because -1 is a square in K. However, $\mathrm{Gal}(h) = SS_5$. We obtained the desired contradiction. $\square$

Example 6.4. Let k be an algebraically closed field of characteristic 3. Let $K = k(z)$ be the field of rational functions in variable z with constant field k. We write $\overline{k(z)}$ for an algebraic closure of $k(z)$. Let $h(x) \in k[x]$ be a *Morse polynomial* of degree 5. This means that the derivative $h'(x)$ of $h(x)$ has $\deg(h) - 1 = 4$ distinct roots $\beta_1, \cdots \beta_4$ and $h(\beta_i) \neq h(\beta_j)$ while $i \neq j$. (For example, $x^5 - x$ is a Morse polynomial.) Then a theorem of Hilbert ([20, theorem 4.4.5, p. 41]) asserts that the Galois group of $h(x) - z$ over $k(z)$ is $\mathbf{S}_n$. Let us consider the hyperelliptic curve

$$C : y^2 = h(x)$$

of genus 2 over $\overline{k(z)}$ and its jacobian $J(C)$. It follows from Theorem 6.3 that $\mathrm{End}(J(C_f)) = \mathbf{Z}$. (The case of $h(x) = x^5 - x$ was earlier treated by Mori [15].)

7. A COROLLARY

Combining Theorems 2.1 and 6.3 together with Theorem 2.3 of [28] and Theorem 2.1 of [25], we obtain the following statement.

Theorem 7.1. *Let K be a field with* $\mathrm{char}(K) \neq 2$, *K_a its algebraic closure, $f(x) \in K[x]$ an irreducible separable polynomial of degree $n \geq 5$ such that the Galois group of f is either $\mathbf{S}_n$ or $\mathbf{A}_n$. If* $\mathrm{char}(K) = 3$ *and $n \leq 6$ then we additionally assume that* $\mathrm{Gal}(f) = \mathbf{S}_n$ *and K contains $\mathbf{F}_9$.*

Let C_f be the hyperelliptic curve $y^2 = f(x)$. Let $J(C_f)$ be its jacobian, $\mathrm{End}(J(C_f))$ *the ring of K_a-endomorphisms of $J(C_f)$. Then* $\mathrm{End}(J(C_f)) = \mathbf{Z}$.

References

[1] S. S. Abhyankar, *Galois theory on the line in nonzero characteristic.* Bull. AMS **27** (1992), 68–133.

[2] L. Dornhoff, Group Representation Theory, Part A, Marcel Dekker, Inc., New York, 1972.

[3] W. Feit, *The computations of some Schur indices.* Israel J. Math. **46** (1983), 274–300.

[4] D. Gorenstein, Finite Simple Groups, An Introduction to their Classification, Plenum Press, New York and London, 1982.

[5] T. Ibukiyama, T. Katsura, F. Oort, *Supersingular curves of genus two and class numbers.* Compositio Math. **57** (1986), 127–152.

[6] I. M. Isaacs, Character theory of finite groups, Pure and Applied Mathematics, No. 69, Academic Press, New York–San Francisco–London, 1976.

[7] A. A. Ivanov, Ch. E. Praeger, *On finite affine 2-Arc transitive graphs.* Europ. J. Combinatorics **14** (1993), 421–444.

[8] G. Janusz, *Simple components of* $\mathbf{Q}[\mathrm{SL}(2,q)]$. Commun. Algebra **1** (1974), 1–22.

[9] N. Katz, *Monodromy of families of curves: applications of some results of Davenport-Lewis.* In: Séminaire de Théorie des Nombres, Paris 1979-80 (ed. M.-J. Bertin); Progress in Math. **12**, pp. 171–195, Birkhäuser, Boston-Basel-Stuttgart, 1981.

[10] N. Katz, *Affine cohomological transforms, perversity, and monodromy.* J. Amer. Math. Soc. **6** (1993), 149–222.

[11] N. Katz, P. Sarnak, Random matrices, Frobenius eigenvalues and Monodromy. AMS, Providence, RI, 1999.

[12] M. Klemm, *Über die Reduktion von Permutationsmoduln.* Math. Z. **143** (1975), 113–117.

[13] Yu. I. Manin, *The theory of commutative formal groups over fields of finite characteristic.* Russian Math. Surveys **18** (1963), 1–83.

[14] D. Masser, *Specialization of some hyperelliptic jacobians.* In: Number Theory in Progress (eds. K. Györy, H. Iwaniec, J.Urbanowicz), vol. I, pp. 293–307; de Gruyter, Berlin-New York, 1999.

[15] Sh. Mori, *The endomorphism rings of some abelian varieties.* Japanese J. Math, **2** (1976), 109–130.

[16] Sh. Mori, *The endomorphism rings of some abelian varieties.*II. Japanese J. Math, **3** (1977), 105–109.

[17] B. Mortimer, *The modular permutation representations of the known doubly transitive groups.* Proc. London Math. Soc. (3) **41** (1980), 1–20.

[18] D. Mumford, Abelian varieties, Second edition, Oxford University Press, London, 1974.

[19] N. O. Nygaard, *Slopes of powers of Frobenius on crystalline cohomology.* Ann. Sci. Ecole Norm. Sup. (4) **14** (1981), no. 4, 369–401.

[20] J.-P. Serre, Topics in Galois Theory. Jones and Bartlett Publishers, Boston-London, 1992.

[21] J.-P. Serre, *Revêtements des courbes algébriques.* Séminaire Bourbaki 1991/92, n 749, Astérisque **206** (1992), 177–182; Œ. vol. IV, 157, 252–264.

[22] A. Silverberg, *Fields of definition for homomorphisms of abelian varieties.* J. Pure Applied Algebra **77** (1992), 253–262.

[23] A. Silverberg, Yu. G. Zarhin, *Variations on a theme of Minkowski and Serre.* J. Pure Applied Algebra **111** (1996), 285–302.

[24] N. Yui, *On the Jacobian varieties of hyperelliptic curves over fields of characteristic* $p > 2$. J. Algebra **52** (1978), 378–410.

[25] Yu. G. Zarhin, *Hyperelliptic jacobians without complex multiplication.* Math. Res. Letters **7** (2000), 123–132.

[26] Yu. G. Zarhin, *Hyperelliptic Jacobians without complex multiplication in positive characteristic.* Math. Research Letters **8** (2001), 429–435.

[27] Yu. G. Zarhin, *Hyperelliptic jacobians and modular representations.* In: "Moduli of abelian varieties" (eds. G. van der Geer, C. Faber, F. Oort); Progress in Math. **195**, pp. 473–490, Birkhäuser, Basel–Boston–Berlin, 2001.

[28] Yu. G. Zarhin, *Very simple* 2*-adic representations and hyperelliptic jacobians.* Moscow Math. J. **2** (2002), issue 2, 403-431.

[29] Yu. G. Zarhin, *Hyperelliptic Jacobians without Complex Multiplication, Doubly Transitive Permutation Groups and Projective Representations.* In: Algebraic Number Theory and Algebraic Geometry (Parshin Festschrift), Contemporary Math. **300** (2002), 195–210.

[30] Yu. G. Zarhin, *Hyperelliptic jacobians and simple groups* $\mathrm{U}_3(2^m)$. Proc. AMS **131** (2003), no. 1, 95–102.

[31] Yu. G. Zarhin, *Very simple representations: variations on a theme of Clifford*. In: Galois Theory Conference Proceedings (H. Völklein, ed.), Developments in Math., Kluwers, to appear; http://front.math.ucdavis.edu/math.GR/0209083 .

Department of Mathematics, Pennsylvania State University, University Park, PA 16802, USA
E-mail address: `zarhin@math.psu.edu`